Maternal Health: Antenatal Care





2014-15 Tanzania Service Provision Assessment (TSPA)

- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare

Antenatal Care by Skilled Provider (2010 TDHS)



^{*}Skilled provider includes doctor/AMO, clinical officer, assistant clinical officer, nurse/midwife, and MCH aide.

Timing of Antenatal Care and Number of Visits (2010 TDHS)

- 15% of women went to their first ANC visit during the 1st trimester of pregnancy, as recommended.
- 43% of women had 4 or more ANC visits, as recommended.
- 2% of women had no ANC visits.

Components of Antenatal Care (2010 TDHS)

- 59% took iron tablets or syrup during last pregnancy
- 68% took anti-malaria drugs
- 53% were informed of signs of pregnancy complications
- 68% had blood pressure measured
- 77% had blood samples taken
- 52% had urine samples taken

Tetanus Toxoid Injections (2010 TDHS)

88% of last births were protected against neonatal tetanus. 48% of women received two or more TT injections during their last pregnancy.


- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare

Availability of Antenatal Care by Facility Type

Percent of all facilities (N=1,188) offering ANC




Availability of Antenatal Care by Managing Authority

Percent of all facilities (N=1,188) offering ANC



Frequency of ANC Services

Among facilities offering ANC services (N=1,005), percent of facilities where ANC services are offered for the indicated number of days per week




- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare

Guidelines and Basic Equipment for ANC Services

Percent of facilities offering ANC services (N=1,005) that have:




Infection Control

Percent of facilities offering ANC services (N=1,005) that have:




Diagnostic Capacity for ANC Tests

Percent of facilities offering ANC services (N=1,005) that have the capacity to conduct the indicated tests in the facility



Availability of Medicines for Routine ANC

Percent of facilities offering ANC services (N=1,005) that have essential medicines and tetanus toxoid vaccine for ANC observed to be available on the day of the survey



- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy




© 2013 Jennifer Applegate, Courtesy of Photoshare

Training and Supervision of ANC Providers


Percent of interviewed providers of ANC services (N=4,252) that received:

- In-service training related to ANC during past 24 months
- Personal supervision during past 6 months



Training of ANC Service Providers

Percent of interviewed ANC service providers (N=4,252) that received specific training



- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare

Characteristics of ANC Clients

Among observed ANC clients (N=4,007)


• Follow-up visit: 54	%
-----------------------	---

- First visit for this pregnancy: 46%
- First pregnancy: 25%

4% of all observed ANC clients were in their
1st trimester of pregnancy (<13 weeks); 51% were in their 3rd trimester (27-42 weeks)


Elements of Client Consultation (First-Visit ANC Clients)

Percent of observed first-visit ANC consultations (N=1,853) that assessed:




Basic Physical Examinations

Percent of all observed ANC consultations (N=4,007) that assessed:




Preventive Interventions

Percent of all observed ANC consultations (N=4,007) that:



Counselling on Pregnancy Complications

Percent of all observed ANC consultations (N=4,007) that included counselling on topics related to pregnancy risk symptoms



Counselling Topics Discussed during ANC Visits

Percent of all observed ANC consultations (N=4,007) that included counselling on the following




- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare


Availability of PMTCT Services by Facility Type

Percent of facilities offering ANC (N=1,005) that provide PMTCT services




PMTCT Services

Percent of facilities offering ANC and any PMTCT services (N=947)



Guidelines, Equipment, Diagnostics, and Medicines for PMTCT Services

Percent of facilities offering ANC and any PMTCT services (N=947)




- Background of Antenatal Care (ANC) services
- Availability of ANC services
- Service Readiness
- Management Practices and Training
- Observed ANC Consultations
- Prevention of Mother-to-Child Transmission of HIV
- Malaria in Pregnancy



© 2013 Jennifer Applegate, Courtesy of Photoshare


Guidelines, Equipment, Diagnostics, and Medicines for Malaria in ANC Facilities

Percent facilities offering ANC (N=1,005)




Malaria-related Interventions during ANC Visits

Percent of all observed ANC consultations (N=4,007)



Malaria-related Training of ANC Providers

Percent of interviewed ANC service providers (N=4,251) that received malaria specific training



Key Findings

- 85% of facilities offer ANC services
- 8% of first-visit ANC client consultations had all elements of client history assessed
- 67% of ANC client consultations discussed any risk symptoms for pregnancy complications
- 52% of facilities provided all nine PMTCT services
- 44% of observed ANC clients were prescribed or given a dose of SP by the provider

