

Tanzania



Reproductive and Child Health Survey

1999

National Bureau of Statistics

MEASURE *DHS+*
Macro International Inc.

Tanzania Reproductive and Child Health Survey 1999

National Bureau of Statistics
Dar es Salaam, Tanzania

Macro International Inc.
Calverton, Maryland, U.S.A.

November 2000



National Bureau of Statistics



United Nations Children's Fund



U.S. Agency for
International Development



United Nations Population Fund



This report presents results from the 1999 Tanzania Reproductive and Child Health Survey (TRCHS) which was undertaken by the National Bureau of Statistics in collaboration with the Reproductive and Child Health Section of the Ministry of Health. Financial assistance for the survey was provided by the U.S. Agency for International Development (USAID/Tanzania), UNICEF/Tanzania, and the United Nations Population Fund (UNFPA/Tanzania). The TRCHS is part of the worldwide MEASURE Demographic and Health Surveys (DHS+) project which is designed to collect, analyse and disseminate data on fertility, family planning, maternal and child health, and HIV/AIDS.

Additional information about the TRCHS may be obtained free of charge from the National Bureau of Statistics, P.O. Box 796, Dar es Salaam (telephone: 135-602; fax: 135-601). Information about the MEASURE DHS+ project may be obtained from Macro International Inc., 11785 Beltsville Drive, Suite 300, Calverton, MD 20705 (telephone: 301-572-0200; fax: 301-572-0999).

Suggested citation:

National Bureau of Statistics [Tanzania] and Macro International Inc. 2000. *Tanzania Reproductive and Child Health Survey 1999*. Calverton, Maryland: National Bureau of Statistics and Macro International Inc.

CONTENTS

	Page
Tables and Figures	vii
Foreword	xiii
Summary of Findings	xv
Map of Tanzania	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Geography, History, and the Economy	1
1.2 Demographic Statistics	2
1.3 Population and Family Planning Policies and Programmes	2
1.4 Health Priorities and Programmes	4
1.5 Objectives and Organisation of the Survey	4
CHAPTER 2 CHARACTERISTICS OF HOUSEHOLDS AND RESPONDENTS	9
2.1 Population by Age and Sex	9
2.2 Household Composition	10
2.3 Fosterhood and Orphanhood	11
2.4 Education Level of Household Population	12
2.5 School Attendance	14
2.6 Child Labour	15
2.7 Housing Characteristics	16
2.8 Background Characteristics of Respondents	19
2.9 Educational Level of Survey Respondents	21
2.10 Access to Media	25
2.11 Employment and Occupation	26
CHAPTER 3 FERTILITY	31
3.1 Current Fertility	31
3.2 Fertility Differentials	32
3.3 Fertility Trends	33
3.4 Children Ever Born	34
3.5 Births Intervals	35
3.6 Age at First Birth	37
3.7 Teenage Pregnancy and Motherhood	37

CHAPTER 4	FERTILITY REGULATION	41
4.1	Knowledge of Family Planning Methods	41
4.2	Trends in Contraceptive Knowledge	43
4.3	Ever Use of Family Planning Methods	44
4.4	Current Use of Family Planning Methods	45
4.5	Sources of Family Planning Methods	51
4.6	Contact of Nonusers with Family Planning Providers	52
4.7	Intention to Use Family Planning Among Nonusers	54
4.8	Reasons for Nonuse	55
4.9	Exposure to Family Planning Messages	55
4.10	Exposure to Reproductive and Child Health Dramas	56
4.11	Knowledge of Family Planning Logo	58
4.12	Knowledge of Salama Condom	62
4.13	Attitudes towards Family Planning	62
CHAPTER 5	PROXIMATE DETERMINANTS OF FERTILITY	65
5.1	Current Marital Status	65
5.2	Age at First Marriage	65
5.3	Age at First Intercourse	67
5.4	Recent Sexual Activity	69
5.5	Postpartum Amenorrhoea, Abstinence, and Insusceptibility	73
5.6	Termination of Exposure to Pregnancy	74
CHAPTER 6	FERTILITY PREFERENCES	75
6.1	Reproductive Preferences	75
6.2	Need for Family Planning Services	78
6.3	Ideal Number of Children	80
6.4	Fertility Planning	83
CHAPTER 7	INFANT AND CHILD MORTALITY	85
7.1	Definitions, Methodology, and Assessment of Data Quality	85
7.2	Childhood Mortality Levels and Trends	86
7.3	Childhood Mortality Differentials	87
7.4	High-Risk Fertility Behaviour	90

CHAPTER 8	REPRODUCTIVE AND CHILD HEALTH	93
8.1	Antenatal Care	93
8.2	Delivery Care	98
8.3	Postnatal Care	102
8.4	Birth Registration	103
8.5	Childhood Vaccinations	104
8.6	Acute Respiratory Infection and Fever	107
8.7	Use of Bednets	108
8.8	Prevalence and Treatment of Diarrhoea	110
8.9	Knowledge of Signs of Illness	111
CHAPTER 9	INFANT FEEDING AND CHILDHOOD NUTRITION	115
9.1	Breastfeeding and Supplementation	115
9.2	Nutritional Status in Early Childhood	119
9.3	Vitamin A Supplementation among Children	123
CHAPTER 10	KNOWLEDGE OF AIDS	125
10.1	Introduction	125
10.2	Knowledge of AIDS Transmission	125
10.3	Perceptions of Personal Risk of Getting AIDS	128
10.4	Knowledge and Use of Condoms	130
10.5	Acceptability of Condom Use	132
10.6	Number of Sexual Partners	133
10.7	AIDS Testing	137
10.8	Knowledge of Condom's Dual Protection	138
REFERENCES		145
APPENDIX A	SAMPLE IMPLEMENTATION	147
APPENDIX B	ESTIMATES OF SAMPLING ERROR	149
APPENDIX C	DATA QUALITY TABLES	165
APPENDIX D	1999 TRCHS HOUSEHOLD SURVEY STAFF	171
APPENDIX E	QUESTIONNAIRES	173

TABLE AND FIGURES

	Page
CHAPTER 1 INTRODUCTION	
Table 1.1 Demographic characteristics	2
Table 1.2 Results of the household and individual interviews	7
CHAPTER 2 CHARACTERISTICS OF HOUSEHOLDS AND RESPONDENTS	
Table 2.1 Household population by age, residence, and sex	9
Table 2.2 Population by age from selected sources	11
Table 2.3 Household composition	11
Table 2.4 Fosterhood and orphanhood	12
Table 2.5.1 Educational level of the female household population	13
Table 2.5.2 Educational level of the male household population	14
Table 2.6 Early childhood education	16
Table 2.7 School attendance ratios	17
Table 2.8 Child labour	18
Table 2.9 Housing characteristics	19
Table 2.10 Household durable goods	20
Table 2.11 Iodised salt	20
Table 2.12 Background characteristics of children under five	21
Table 2.13 Background characteristics of respondents	22
Table 2.14.1 Level of education: women	23
Table 2.14.2 Level of education: men	23
Table 2.15 Literacy	24
Table 2.16 Access to mass media	25
Table 2.17 Employment	27
Table 2.18 Employer and form of earnings	28
Table 2.19.1 Occupation: women	29
Table 2.19.2 Occupation: men	30
Figure 2.1 Population pyramid, Tanzania 1999	10
Figure 2.2 Level of education for women and men, Tanzania 1999	15
Figure 2.3 Percentage of children age 7-13 attending primary school, by sex and residence	17
CHAPTER 3 FERTILITY	
Table 3.1 Current fertility	31
Table 3.2 Fertility by background characteristics	33
Table 3.3 Trends in fertility	34
Table 3.4 Age-specific fertility rates	34

Table 3.5	Children ever born and living	35
Table 3.6	Birth intervals	36
Table 3.7	Age at first birth	37
Table 3.8	Median age at first birth by background characteristics	38
Table 3.9	Teenage pregnancy and motherhood	38
Figure 3.1	Age-specific fertility rates by urban-rural residence	32
Figure 3.2	Total fertility rates by background characteristics	33
Figure 3.3	Pregnancy and childbearing among women age 15-19	39

CHAPTER 4 FERTILITY REGULATION

Table 4.1	Knowledge of contraceptive methods	42
Table 4.2	Couples' knowledge of contraceptive methods	43
Table 4.3.1	Ever use of contraception: women	45
Table 4.3.2	Ever use of contraception: men	46
Table 4.4.1	Current use of contraception: women	47
Table 4.4.2	Current use of contraception: men	48
Table 4.5.1	Current use of contraception by background characteristics: women	49
Table 4.5.2	Current use of contraception by background characteristics: men	51
Table 4.6	Source of supply for contraceptive methods	53
Table 4.7	Contact of nonusers with family planning providers	54
Table 4.8	Future use of contraception	55
Table 4.9	Reasons for not intending to use contraception	56
Table 4.10.1	Heard family planning message: women	57
Table 4.10.2	Heard family planning message: men	58
Table 4.11	Exposure to family planning dramas	59
Table 4.12.1	Green Star logo family planning symbol: women	60
Table 4.12.2	Green Star logo family planning symbol: men	61
Table 4.13	Knowledge of Salama condom	62
Table 4.14	Discussion of family planning with husband	63
Table 4.15	Attitudes of couples toward family planning	64
Table 4.16	Spouse's perception of their spouse's approval of family planning	64
Figure 4.1	Trends in contraceptive knowledge among women age 15-49, 1991-1999	44
Figure 4.2	Contraceptive use among all women age 15-49 by residence and education	50
Figure 4.3	Trends in contraceptive use among all women age 15-49, 1991-1999	52

CHAPTER 5 PROXIMATE DETERMINANTS OF FERTILITY

Table 5.1	Current marital status	66
Table 5.2	Age at first marriage	67
Table 5.3	Median age at first marriage	68
Table 5.4	Age at first sexual intercourse	69
Table 5.5	Median age at first intercourse	70
Table 5.6.1	Recent sexual activity: women	71
Table 5.6.2	Recent sexual activity: men	72
Table 5.7	Postpartum amenorrhoea, abstinence, and insusceptibility	73
Table 5.8	Median duration of postpartum insusceptibility by background characteristics	74
Table 5.9	Menopause	74

CHAPTER 6 FERTILITY PREFERENCES

Table 6.1	Fertility preferences by number of living children	75
Table 6.2	Fertility preferences by age	77
Table 6.3	Desire to stop childbearing by background characteristics	78
Table 6.4	Need for family planning	79
Table 6.5	Ideal number of children	81
Table 6.6	Mean ideal number of children	82
Table 6.7	Fertility planning status	83
Table 6.8	Wanted fertility rates	84
Figure 6.1	Fertility preferences of women age 15-49	76
Figure 6.2	Desire to stop childbearing among women age 15-49, by number of living children	77
Figure 6.3	Trends in mean ideal family size among women and men, 1991-1999	82

CHAPTER 7 INFANT AND CHILD MORTALITY

Table 7.1	Infant and child mortality	86
Table 7.2	Infant and child mortality by background characteristics	87
Table 7.3	Infant and child mortality by biodemographic characteristics	89
Table 7.4	High-risk fertility behaviour	91
Figure 7.1	Under-five mortality by selected background characteristics	88

CHAPTER 8 REPRODUCTIVE AND CHILD HEALTH

Table 8.1	Antenatal care	94
Table 8.2	Number of antenatal care visits and stage of pregnancy	95
Table 8.3	Antenatal care content	96

Table 8.4	Tetanus toxoid vaccinations	97
Table 8.5	Place of delivery	99
Table 8.6	Assistance during delivery	100
Table 8.7	Delivery characteristics	101
Table 8.8	Postnatal care	102
Table 8.9	Postpartum vitamin A supplements	103
Table 8.10	Birth registration coverage	104
Table 8.11	Vaccinations by source of information	105
Table 8.12	Vaccinations by background characteristics	107
Table 8.13	Prevalence and treatment of acute respiratory infection and prevalence of fever	109
Table 8.14	Use of bednets	110
Table 8.15	Prevalence of diarrhoea	111
Table 8.16	Treatment of diarrhoea	112
Table 8.17	Feeding practices during diarrhoea	113
Table 8.18	Knowledge of health complications	113
Table 8.19	Knowledge of pregnancy complications	114
Figure 8.1	Antenatal and delivery care indicators	98
Figure 8.2	Percentage of children age 12-23 months with specific vaccinations according to vaccination cards and mothers' reports	106

CHAPTER 9 INFANT FEEDING AND CHILDHOOD NUTRITION

Table 9.1	Breastfeeding status	116
Table 9.2	Median duration and frequency of breastfeeding	117
Table 9.3	Infant feeding indicators	118
Table 9.4	Ideal duration of breastfeeding	119
Table 9.5	Nutritional status of children	122
Table 9.6	Trends in nutritional status of children	123
Table 9.7	Vitamin A supplementation among children	124

CHAPTER 10 KNOWLEDGE OF AIDS

Table 10.1	Knowledge of HIV/AIDS	126
Table 10.2.1	Knowledge of specific ways to avoid HIV/AIDS: women	127
Table 10.2.2	Knowledge of specific ways to avoid HIV/AIDS: men	128
Table 10.3	Knowledge (prompted) of the main ways to avoid HIV/AIDS	129
Table 10.4	Misconceptions about HIV/AIDS transmission	130
Table 10.5.1	Knowledge and perception of HIV/AIDS: women	131
Table 10.5.2	Knowledge and perception of HIV/AIDS: men	132
Table 10.6	Knowledge of mother-child transmission of HIV/AIDS	133
Table 10.7	Perception of the risk of getting HIV/AIDS	134
Table 10.8	Reasons for perception of small/no risk of getting HIV/AIDS	135

Table 10.9	Reasons for perception of moderate/great risk of getting HIV/AIDS	135
Table 10.10	Knowledge of condoms	136
Table 10.11.1	Use of condoms: women	137
Table 10.11.2	Use of condoms: men	138
Table 10.12	Acceptability of women prompting use of condoms	139
Table 10.13.1	Number of sexual partners: women	140
Table 10.13.2	Number of sexual partners: men	141
Table 10.14.1	Testing for HIV/AIDS: women	142
Table 10.14.2	Testing for HIV/AIDS: men	143
Table 10.15	Knowledge of dual protection of condoms	144

APPENDIX A SAMPLE IMPLEMENTATION

Table A.1.1	Sample implementation: women	147
Table A.1.2	Sample implementation: men	148

APPENDIX B ESTIMATES OF SAMPLING ERROR

Table B.1	List of selected variables for sampling errors, Tanzania 1999	152
Table B.2	Sampling errors for women - Total sample: Tanzania 1999	153
Table B.3	Sampling errors for women - Urban sample: Tanzania 1999	154
Table B.4	Sampling errors for women - Rural sample: Tanzania 1999	155
Table B.5	Sampling errors for women - Mainland sample: Tanzania 1999	156
Table B.6	Sampling errors for women - Zanzibar sample: Tanzania 1999	157
Table B.7	Sampling errors for women - Pemba sample: Tanzania 1999	158
Table B.8	Sampling errors for women - Unguja sample: Tanzania 1999	159
Table B.9	Sampling errors for men - Total sample: Tanzania 1999	160
Table B.10	Sampling errors for men - Urban sample: Tanzania 1999	160
Table B.11	Sampling errors for men - Rural sample: Tanzania 1999	161
Table B.12	Sampling errors for men - Mainland sample: Tanzania 1999	161
Table B.13	Sampling errors for men - Zanzibar sample: Tanzania 1999	162
Table B.14	Sampling errors for men - Pemba sample: Tanzania 1999	162
Table B.15	Sampling errors for men - Unguja sample: Tanzania 1999	163

APPENDIX C DATA QUALITY TABLES

Table C.1	Household age distribution	165
Table C.2	Age distribution of eligible and interviewed women	166
Table C.3	Completeness of reporting	166
Table C.4	Births by calendar years	167
Table C.5	Reporting of age at death in days	168
Table C.6	Reporting of age at death in months	169

FOREWORD

The National Bureau of Statistics takes pleasure in presenting this report on the 1999 Reproductive and Child Health Survey (TRCHS). The 1999 TRCHS is the latest in a series of periodic surveys to measure levels, patterns, and trends in demographic and health indicators, the first having been the 1991-92 Tanzania Demographic and Health Survey (TDHS).

This report contains findings from the 1999 TRCHS regarding data from the households visited. The tables and text cover the most important indicators and should be of use to policy makers and program administrators who need up-to-date data for evaluating their activities and planning future directions. Findings from the survey covering health facilities will be produced in a separate report.

The successful completion of the 1999 TRCHS was made possible by the joint effort of a number of organisations and individuals, whose participation we would like to acknowledge with gratitude. First, is the U.S. Agency for International Development (USAID)/Tanzania which has long supported the collection and utilisation of data to evaluate the family planning and health programmes in Tanzania and which initiated planning for this survey and provided the bulk of the funding to implement it. UNICEF/Tanzania and UNFPA/Tanzania also contributed substantially to both the survey design and the funding for the survey. Many other organisations contributed to the questionnaire content and/or the field staff training, including the Reproductive and Child Health Unit at the Ministry of Health, the Tanzania Food and Nutrition Centre, and the National AIDS Control Programme and Mount Meru Hospital. We would also like to thank the Demographic and Health Surveys program of Macro International Inc. in Calverton, Maryland, U.S.A. for providing technical assistance in all phases of the project. The survey would not have gotten off the ground without the exemplary and tireless efforts of the staff at the National Bureau of Statistics. Their many long days of overtime work have served to make this survey effort a success. Similarly, the nurses who acted as interviewers for the survey deserve our heartfelt thanks. Finally, we are ever more grateful to the survey respondents who contributed generously part of their time to enable us to gather crucial data for our country's future planning.

Cletus P.B. Mkai
Director General
National Bureau of Statistics

Dar es Salaam, Tanzania

SUMMARY OF FINDINGS

The 1999 Tanzania Reproductive and Child Health Survey (TRCHS) is a nationally-representative sample survey covering 4,029 women age 15-49 and 3,542 men age 15-59. The TRCHS was designed to provide information on levels and trends of fertility, family planning knowledge and use, infant and child mortality, and indicators of maternal and child health and nutrition. Fieldwork for the TRCHS took place from early September to late-November 1999.

Survey data generally confirm patterns observed in the 1996 Tanzania Demographic and Health Survey (TDHS), showing increasing contraceptive use and more widespread knowledge about HIV/AIDS; however, results show that many challenges still exist.

FERTILITY

Fertility Decline. The TRCHS data indicate that there has been a small decline in fertility since the 1996 survey. The total fertility rate has dropped from 5.8 births per woman in the period 1994-96 to 5.6 births for the period 1997-99. The rate of decline may be slowing somewhat, however, and the level of fertility is still high.

Large Fertility Differentials. Significant differences in fertility levels are evident from survey data. For example, the total fertility rate among rural women is twice that of urban women (6.5 versus 3.2). Fertility levels are closely related to women's education. Women with no formal education give birth to an average of 6.5 children in their lifetime, compared with 4.9 for women who have completed primary school.

Unplanned Fertility. One reason for the relatively high fertility levels is that unplanned pregnancies are still common. Overall, more than one in five births in the three years prior to the survey were reported to be unplanned; 11 percent were mistimed (wanted

later) and 11 percent were unwanted. Unwanted births are disproportionately high among older women who already have several children, the very women who are at higher risk of fertility-related illness and higher child mortality. If unwanted births could be eliminated altogether, the total fertility rate in Tanzania would be 4.8 births per woman instead of the actual level of 5.6.

Ideal Family Size. Although a reduction in the number of unplanned births would reduce fertility substantially, Tanzanian women and men still want to have large families. Even those who have two children or fewer say they would ideally like to have almost five children on average.

FAMILY PLANNING

Increasing Use of Contraception. A major cause of the declining fertility in Tanzania has been the slow but steady increase in contraceptive use over the last decade. The contraceptive prevalence rate has doubled since 1991-92, from 10 to 22 percent of all women. Use of modern methods has grown from 6 to 16 percent of all women. Overall, there has been a steady growth in the contraceptive prevalence rate with an average increase of one and a half percentage points a year.

Method Mix. In terms of "method mix," the dominant change over the last 8 years has been the large increase in the number of women using injectable contraception. The proportion of women relying on injectables increased from less than half a percent in 1991-92 to over 5 percent in 1999 and the injectable now accounts for one-quarter of all contraceptive use.

Use of condoms, periodic abstinence and withdrawal have also increased somewhat over the past few years. A levelling off in pill use may indicate that some women are switch-

ing from the pill to injectables. The low level of use of permanent methods such as sterilisation and implants is of concern, given the high level of unplanned births among high-parity, older women.

Differentials in Family Planning Use.

Differentials in current use of family planning are large. Urban women are almost twice as likely as rural women to be using a contraceptive method (33 versus 18 percent). Women in the Mainland are more likely to use than those on Zanzibar. However, the largest differences are found by education—contraceptive use among women with some secondary education is three times higher than among those with no education.

Knowledge of Contraception. The proportion of women and men who know of at least one contraceptive method has been over 75 percent for some time and the TRCHS results indicate that the proportion has increased to over 90 percent. Moreover, the mean number of methods that women and men say they have heard of has grown from 4 in 1996 to 6 in 1999. Knowing about more methods provides a basis for wider choice.

Unmet Need for Family Planning.

Unmet need for family planning has declined slightly since 1996. Data from the 1996 TDHS show that 19 percent of all women were in need of services, compared with 17 percent in the 1999 TRCHS. Two-thirds of the unmet need is comprised of women who want to space their next birth, while just one third is for women who do not want any more children (limiters). If all women who say they want to space or limit their children were to use methods, the contraceptive prevalence rate could be increased from 22 percent to 40 percent of all women. Currently, 56 percent of this “total demand” for family planning is being met.

Family Planning Messages. Survey data show that family planning messages are continuing to get through to couples. As in 1996, just over 40 percent of women have

heard a family planning message on the radio in the six months prior to the survey, while 5 percent have seen a message on the television and one-fifth have seen a poster. Taking all sources into account, two-thirds of women and 70 percent of men say they have seen or heard a family planning message in the last six months. The proportion of women and men who say they have seen or heard specific programs such as *Zinduka!* and *Twende na Wakati* has also increased since 1996.

MATERNAL AND CHILD HEALTH

Maternal Health Care. TRCHS data point to several areas regarding maternal health care in which improvements could be made. Although the proportion of Tanzanian mothers who receive antenatal care from a doctor, nurse, midwife or medical aide has remained steady at just over 90 percent since 1991-92, there has been a shift in providers from nurses and midwives to the less well-trained health aides. The proportion of pregnant women receiving at least one tetanus toxoid injection has declined from 92 percent in 1996 to 83 in 1999. The data also show a disturbingly steady decline in the proportion of births that occur in a health facility—from 53 percent in 1991-92 to 47 percent in 1996 to 44 percent in 1999. Because of this decline, the proportion of births assisted by trained medical personnel (doctors, nurses, midwives) has declined from 44 percent in 1991-92 to 36 percent in 1999. Less than one in five of those who deliver at home go to a health facility for a postnatal check-up within a month after delivery.

Possible Leveling Off of Childhood Mortality Decline. Survey results imply that the decline in childhood mortality documented in the 1996 TDHS may be stagnating or even increasing slightly. The TRCHS rates show that almost 1 in 7 children born in Tanzania dies before reaching the fifth birthday, an indication that there is still much improvement to be made. The under-five mortality rate measured in the survey is 147 deaths per 1,000 births; the infant mortality rate is 99 per 1,000.

Childhood Vaccination Coverage. The 1999 TRCHS results show that 68 percent of children age 12-23 months are fully vaccinated, close to the 71 percent in 1996.

Childhood Health. The TRCHS provides data on some of the more common childhood illness and their treatment. Just over 1 in 3 children under age five had a fever and 14 percent had respiratory illness in the two weeks before the survey. Of these, two-thirds were taken to a health facility for treatment. Twelve percent of children under five were reported to have had diarrhoea in the two weeks preceding the survey. The fact that two-thirds of children with diarrhoea received some type of oral rehydration therapy (fluid made from an ORS packet or increased fluids) is encouraging.

Breastfeeding Practices. The TRCHS results document a relatively long duration of breastfeeding, with a median duration of 21 months. Although breastfeeding has beneficial effects on both the child and the mother, TRCHS data indicate that supplementation of breastfeeding with other liquids and foods occurs too early in Tanzania. For example, among newborns less than four months of age, 60 percent are already receiving complementary foods or liquids.

Nutritional Status of Children. Results show no appreciable change in the nutritional status of children in Tanzania. Over 40 percent of children under five show evidence of chronic malnutrition or stunting, while 5 percent are acutely malnourished (wasted).

Vitamin A Supplementation. Over the past decade, several studies have proved the importance of adequate vitamin A in mitigating the severity of maternal and childhood illnesses and thereby reducing mortality. Supplementing young children and postpartum women with a capsule containing a high dose of vitamin A is an easy way to ensure adequate intake. However, survey data show that only 14 percent of children aged 6-59 months received a high-dose vitamin A

supplement in the six months prior to the survey. Vitamin A supplementation for postpartum women has about the same coverage; 12 percent of women received a supplement within 2 months after delivering.

HIV/AIDS-RELATED INDICATORS

Awareness of AIDS. The TRCHS results confirm the fact that 97 percent of women and 99 percent of men have heard of HIV/AIDS. They also indicate an improvement in knowledge about how the disease is spread, which is reflected in an increase in the proportion of women and men who can list most of the major ways to avoid getting the illness. Around three-quarters or more of respondents know that HIV can be passed from a mother to her child and that a healthy-looking person can be infected.

Condom Use. One of the main objectives of the Tanzanian AIDS control programme is to encourage consistent use of condoms, especially with partners who may be exposed to HIV with other partners or through other risky behaviour. TRCHS data show that condom use with extramarital partners has increased since 1996 for women but not for men. Almost one in four women and 34 percent of men who had sex with someone other than a husband or wife in the previous 12 months said they used a condom the last time. Half of the respondents feel it is acceptable for a woman to ask a man to use a condom, while almost 60 percent say that if a man has a sexually transmitted disease, it is acceptable for his wife to ask him to use condoms or to refuse to have sex with him.

HIV Testing. TRCHS data show that 7 percent of women and 12 percent of men have been tested for HIV, which represents a slight increase from 1996 (4 and 11 percent, respectively). As before, about two-thirds of those not tested say they would like to be. Major reasons for not getting tested are not knowing where to go and not having the time to go.

TANZANIA

